arXiv:1212.0255v1 [math.PR] 2 Dec 2012

Einstein Relation for Random Walks in Balanced Random Environment

Xiaoqin Guo*

December 01, 2012

Abstract

The Einstein relation describes the relation between the response of a system to a perturbation and its diffusivity at equilibrium. It states that the derivative of the velocity (with respect to the strength of the perturbation) equals the diffusivity. This paper consider the Einstein relation in the context of random walks in a balanced uniformly elliptic iid environment. We show that in our context, the Einstein relation is a consequence of the convergence of stationary measures. Our argument is based on a new regeneration structure for the random walks in the perturbed environment.

1 Introduction

In the 1900s, Einstein [4, pp. 1-18] investigated the movement of suspended particles in a liquid under the influence of an external force. He established the following linear relation between the diffusion constant D and the *mobility* (i.e. the ratio between the effective velocity and the force) μ:

$$D \sim \mu.$$

Recently, there has been much interest in studying the Einstein relation for reversible motions in random media, see [13, 8, 6, 1]. So far, little work has been done in the non-reversible zero speed set-up, e.g. random walks in random environments (RWRE). The purpose of this article is to prove the Einstein relation for random walks in a perturbation of a zero-drift random environment, i.e, the derivative of the velocity with respect to the size of the perturbation converges to the diffusivity constant of the environment.

In this section, we will first define the model of RWRE and then formulate the problem.

---

*Technische Universität München, Lehrstuhl für Wahrscheinlichkeitstheorie, 85748 Garching, Germany. Research supported by NSF grant DMS-0804133.

An *environment* $\omega : \mathbb{Z}^d \times \{e \in \mathbb{Z}^d : |e| = 1\} \to [0,1]$ is a function that satisfies

$$\sum_{e:|e|=1} \omega(x,e) = 1$$

for all $x \in \mathbb{Z}^d$, where $|\cdot|$ denotes the l^2-norm. We endow the set of all environments Ω with a probability measure P. The random environment is called *balanced* if

$$P(\omega(x,e) = \omega(x,-e) \text{ for all } x, e \in \mathbb{Z}^d) = 1,$$

and *uniformly elliptic* if $\omega > \kappa$ for almost every ω and some $\kappa > 0$.

The random walks $(X_n)_{n\geq 0}$ in the random environment ω is defined to be the Markov chain with transition probability P_ω specified by

$$\begin{aligned}&P_\omega^x(X_0 = x) = 1,\\ &P_\omega^x(X_{n+1} = y + e | X_n = y) = \omega(y,e).\end{aligned}$$

The measure P_ω^x of the paths under a fixed environment ω is called the *quenched law.* The average over all quenched environments, $P \otimes P_\omega^x$, is called the *annealed law.* Expectation with respect to P_ω^x is denoted by E_ω^x.

For $\omega \in \Omega$, set

$$\omega_x = \big(\omega(x,e)\big)_{|e|=1}.$$

Define the spatial shifts $\{\theta^y\}_{y\in\mathbb{Z}^d}$ on Ω by $(\theta^y\omega)_x = \omega_{x+y}$. We say that the random environment is *iid* if the probability vectors $(\omega_x)_{x\in\mathbb{Z}^d}$ are independent and identically distributed.

For $t \geq 0$, let

$$X_t = X_{\lfloor t\rfloor} + (t - \lfloor t\rfloor)(X_{\lfloor t\rfloor+1} - X_{\lfloor t\rfloor}).$$

Lawler [11] proved a quenched invariance principle for random walks in an ergodic (ergodic with respect to the spatial shifts), balanced and uniformly elliptic random environment. Namely, for P-almost all ω, the P_ω law of $(\lambda X_{t/\lambda^2})_{t\geq 0}$ converges weakly to a Brownian motion with covariance matrix

$$\mathbb{D} = (2E_Q\omega(o,e_i)\delta_{ij})_{1\leq i,j\leq d}$$

as $\lambda \downarrow 0$, where Q is an ergodic initial measure for the "environment viewed from the point of view of the particle" process

$$\bar{\omega}_n := \theta^{X_n}\omega, \quad n \geq 0.$$

See [7, pages 211-212] for the construction of the measure Q. (For random walks in iid balanced environment, it is shown in [7] that the invariance principle still holds assuming only ellipticity, i.e., $\omega > 0$ for P-almost every

ω. Berger and Deuschel [2] have further generalized the result to the non-elliptic case where the environment is "genuinely d-dimensional".)

Now we are ready to consider the random walks in a perturbation of the balanced environment. Throughout this paper, we assume that *the environment P is iid, balanced, and uniformly elliptic with ellipticity constant $\kappa > 0$.*

For $\lambda \in (0,1)$ and a fixed direction

$$\ell = (\ell_1, \ldots, \ell_d) \in S^{d-1},$$

define the perturbed environment ω^λ of $\omega \in \Omega$ by

$$\omega^\lambda(x,e) = (1 + \lambda \ell \cdot e)\omega(x,e).$$

We denote the annealed probability measure of the perturbed environment by $\mathbb{P}_\lambda := P \otimes P^o_{\omega^\lambda}$ and let $\mathbb{E}_\lambda$ denotes its expectation. Since ω^λ satisfies Kalikow's condition (see (0.7) in [15]), it follows from [15, Theorem 2.3] that there exists a deterministic constant $v_\lambda \in \mathbb{R}^d$ such that

$$\lim_{t\to\infty} \frac{X_t}{t} = v_\lambda, \qquad \mathbb{P}_\lambda\text{-almost surely.}$$

Moreover, it follows by [15, Theorem 3.1] that:

> As $n \to \infty$, $\bar{\omega}_n$ (under the law $\mathbb{P}_\lambda$) converges weakly to an invariance measure Q_λ for the process $(\bar{\omega}_i)_{i\in\mathbb{N}}$.

Notice that under $\mathbb{P}_\lambda$, $(\bar{\omega}_n)_{n\ge 0}$ is the process of *balanced environments viewed from the point of view of particle in the perturbed environment*, and so Q_λ is supported on the set of balanced environments.

The following mobility-diffusivity relation

$$\lim_{\lambda\to 0} \frac{v_\lambda}{\lambda} = \mathbb{D}\ell \tag{ER}$$

is called the *Einstein relation* for the RWRE.

In this paper, we will provide a new interpretation of this relation. Namely, for random walks in balanced random environment, the Einstein relation is just a consequence of *the convergence of the invariance measures Q_λ to the equilibrium measure Q*. Indeed, since the drift of ω^λ at $x \in \mathbb{Z}^d$ is

$$\sum_{e:|e|=1} \omega^\lambda(x,e)e = \lambda \sum_{i=1}^d 2\omega(x,e_i)\ell_i e_i := \lambda D(\theta^x \omega),$$

we get that

$$Y_n := \lambda \sum_{i=1}^n D(\bar{\omega}_{i-1}) - X_n \tag{1}$$

is a P_{ω^λ}-martingale with bounded increments, and so $Y_n/n \to 0$, P_{ω^λ}-almost surely. Therefore, $\mathbb{P}_\lambda$-almost surely,

$$v_\lambda = \lambda \lim_{n\to\infty} \frac{1}{n} \mathbb{E}_\lambda \sum_{i=0}^{n-1} D(\bar\omega_i) = \lambda E_{Q_\lambda} D. \tag{2}$$

Since $\mathbb{D}\ell = (2E_Q\omega(o,e_i)\ell_i)_{1\le i\le d} = E_Q D$, the Einstein relation (ER) is an immediate corollary of the following theorem:

Theorem 1. *Assume that the environment is iid, balanced, and uniformly elliptic and the measures Q and Q_λ are defined as above, then*

$$Q_\lambda \Rightarrow Q \quad \text{as } \lambda \to 0,$$

where $\Rightarrow$ denotes weak convergence.

The main purpose of our paper is to prove this theorem.

Since the space Ω of the environment is compact under the product topology, to prove Theorem 1, it suffices to show that $\lim_{\lambda\to 0} E_{Q_\lambda} f = E_Q f$ for all $f : \Omega \to \mathbb{R}$ such that

$$\begin{aligned} &f \text{ is } \sigma(\omega_x : |x| \le N_f)\text{-measurable and } |f| \le A_f \\ &(P\text{-a.s.}) \text{ for some constants } N_f \ge 1 \text{ and } A_f. \end{aligned} \tag{3}$$

Our proof of Theorem 1 consists of proving the following two theorems:

Theorem 2. *Assume that $f : \Omega \mapsto \mathbb{R}$ is $L^1(P)$. Then, for P-almost every ω and any $t > 0$,*

$$\lim_{\lambda\to 0} \frac{\lambda^2}{t} E_{\omega^\lambda}[\sum_{i=0}^{\lceil t/\lambda^2\rceil} f(\bar\omega_i)] = E_Q f.$$

Theorem 3. *Assume the environment P is iid, balanced and uniformly elliptic. Then for any f that satisfies* (3),

$$\left| E_{Q_\lambda} f - \frac{\lambda^2}{t} \mathbb{E}_\lambda \sum_{i=0}^{\lceil t/\lambda^2\rceil} f(\bar\omega_i) \right| \le \frac{C N_f A_f}{t^{1/3}} \quad \forall \lambda \in (0,1), t > 0.$$

Our proof of Theorem 2 is an adaption of the argument of Lebowitz and Rost [13] (see also [6, Proposition 3.1]) to the discrete setting.

For the proof of Theorem 3, we want to follow the strategy of Gantert, Mathieu and Piatnitski [6]. Arguments in the proof of [6, Proposition 5.1] show that if we can construct a sequence of random times $(\tau_n)_{n\in\mathbb{N}}$ (called the *regneration times*) that divides the random path into iid (under the annealed measure) pieces, then good moment estimates of the regeneration times yield the Einstein relation. In [6], a heat kernel estimate [6, Lemma

5.2] for reversible diffusions is crucially employed in the construction of the regeneration times. However, due to the lack of reversibility, we don't have a good heat kernel estimate for RWRE. In this paper, we construct the regeneration times differently, so that they divide the random path into "almost iid" parts. Moreover, our regeneration times have good moment bounds, which lead to a proof of Theorem 3. The key ingredients in our construction are Kuo and Trudinger's [9] Harnack inequality for discrete harmonic functions and the "ϵ-coins" trick introduced in [3].

The structure of the paper is as follows. We will prove Theorem 2 in Section 2. In Section 3, we present our new construction of the regeneration times. Furthermore, we will show in Section 4 that these regeneration times have good moment properties. Section 5 is devoted to the proof of Theorem 3.

Without loss of generality, we assume $\ell_1 = \ell \cdot e_1 > 0$ throughout the paper.

2 Proof of Theorem 2

Proof: We first consider the Radon-Nikodym derivative of the measure P_{ω^λ} with respect to P_ω. For $s > 0$, put

$$G(s,\lambda) = G(s,\lambda; X.) := \log \prod_{j=1}^{\lceil s \rceil} [1 + \lambda \ell \cdot (X_j - X_{j-1})].$$

Then, for any measurable function F on $C([0,s],\mathbb{R}^d)$,

$$E_{\omega^\lambda} F(X_r : 0 \le r \le s) = E_\omega F(X_s : 0 \le r \le s) e^{G(s,\lambda)}.$$

In particular,

$$E_\omega e^{G(s,\lambda)} = E_{\omega^\lambda}[1] = 1 \tag{4}$$

for any $\lambda \in (0,1)$ and $s > 0$. Moreover, by Taylor's expansion, we get

$$\begin{aligned} G(s,\lambda) &= \sum_{j=1}^{\lceil s \rceil} \log(1 + \lambda \ell \cdot (X_j - X_{j-1})) \\ &= \sum_{j=1}^{\lceil s \rceil} \left[\lambda \ell \cdot (X_j - X_{j-1}) - \frac{\lambda^2 \big(\ell \cdot (X_j - X_{j-1})\big)^2}{2} \right] + \lambda^2 \lceil s \rceil H(\lambda) \\ &= \lambda X_{\lceil s \rceil} \cdot \ell - \frac{\lambda^2}{2} \sum_{j=1}^{\lceil s \rceil} \big(\ell \cdot (X_j - X_{j-1})\big)^2 + \lambda^2 \lceil s \rceil H(\lambda), \end{aligned} \tag{5}$$

where the random variable $H(\lambda) = H(\lambda; X.)$ satisfies $0 \leq H \leq \lambda/3$. Setting $h(\omega) = \sum_{i=1}^{d} \omega(o, e_i)\ell_i^2$,

$$\left(\sum_{j=1}^{n} \big(\ell \cdot (X_j - X_{j-1})\big)^2 - 2h(\omega_{X_{j-1}}) \right)_{n \geq 0}$$

is a martingale sequence with bounded increments. Thus P_ω-almost surely,

$$\lim_{n\to\infty} \frac{1}{n} \sum_{j=1}^{n} \left[\big(\ell \cdot (X_j - X_{j-1})\big)^2 - 2h(\theta^{X_{j-1}}\omega) \right] = 0.$$

Further, recall that Q is the ergodic initial measure for the sequence $(\bar{\omega}_n)_{n\geq 0}$ (under the balanced environment ω) and $Q \sim P$. Hence by the ergodic theorem, $\mathbb{P}$-almost surely,

$$\lim_{\lambda\to 0} \lambda^2 \sum_{j=1}^{\lceil t/\lambda^2 \rceil} \big(\ell \cdot (X_j - X_{j-1})\big)^2 = \lim_{\lambda\to 0} \lambda^2 \sum_{j=1}^{\lceil t/\lambda^2 \rceil} 2h(\bar{\omega}_{j-1}) = 2tE_Q h \tag{6}$$

and

$$\lim_{\lambda\to 0} \frac{t}{\lambda^2} \sum_{i=0}^{\lceil t/\lambda^2 \rceil} f(\bar{\omega}_i) = E_Q f. \tag{7}$$

Recall that by the quenched invariance principle [7, Theorem 2] for balanced environments, $(\lambda X_{s/\lambda^2})_{s\geq 0}$ converges weakly (under P_ω) to $(N_s)_{s\geq 0}$. Hence, by (5), (6), (7) and the invariance principle, for P-almost all ω,

$$\frac{t}{\lambda^2} \sum_{i=0}^{\lceil t/\lambda^2 \rceil} f(\bar{\omega}_i) e^{G(t/\lambda^2, \lambda)} \tag{8}$$

converges weakly (under P_ω) to

$$(E_Q f) \exp(N_t \cdot \ell - tE_Q h).$$

Next, we will prove that for P-almost every ω, this convergence is also in $L^1(P_\omega)$. It suffices to show that the class $(e^{G(t/\lambda^2,\lambda)})_{\lambda\in(0,1)}$ is uniformly integrable under P_ω, P-a.s.. Indeed, for any $\gamma > 1$, it follows from (5) and the estimate on $H(\lambda)$ that

$$\begin{aligned}
&\gamma G(t/\lambda^2, \lambda) \\
&\leq G(t/\lambda^2, \gamma\lambda) + \frac{(\gamma^2 - \gamma)\lambda^2}{2} \sum_{j=1}^{\lceil t/\lambda^2 \rceil} \big(\ell \cdot (X_j - X_{j-1})\big)^2 + \gamma\lambda^2 \lceil t/\lambda^2 \rceil H(\lambda) \\
&< G(t/\lambda^2, \gamma\lambda) + \gamma^2(t+1).
\end{aligned}$$

Hence for $\gamma > 1$ and all $\lambda \in (0,1)$,

$$E_\omega \exp(\gamma G(t/\lambda^2, \lambda)) \le e^{\gamma^2(t+1)} E_\omega \exp(G(t/\lambda^2, \gamma\lambda)) \overset{by (4)}{=} e^{\gamma^2(t+1)},$$

which implies the uniform integrability of $(e^{G(t/\lambda^2,\lambda)})_{\lambda\in(0,1)}$. So the $L^1(P_\omega)$-convergence of (8) is proved and (for P-almost every ω)

$$\lim_{\lambda\to 0} E_{\omega^\lambda}\left[\frac{t}{\lambda^2}\sum_{i=0}^{\lceil t/\lambda^2\rceil} f(\bar\omega_i)\right] = (E_Q f)E\big[\exp(N_t\cdot\ell - tE_Q h)\big].$$

The theorem follows by noting that $tE_Q h = E(N_t\cdot\ell)^2/2$ and that

$$E\big[\exp(N_t\cdot\ell - E(N_t\cdot\ell)^2/2)\big] = 1. \quad \square$$

3 Regenerations

3.1 Auxiliary estimates

Let

$$\lambda_1 := 0.5/\lceil(2\lambda\ell_1)^{-1}\rceil$$

so that $0.5/\lambda_1$ is an integer and

$$\frac{1}{2\lambda\ell_1} \le \frac{1}{2\lambda_1} < \frac{1}{2\lambda\ell_1} + 1.$$

For any $n\in\mathbb{Z}, x\in\mathbb{Z}^d$, call

$$\mathcal{H}_n^x = \mathcal{H}_n^x(\lambda,\ell) := \{y\in\mathbb{Z}^d : (y-x)\cdot e_1 = n/\lambda_1\}$$

the n-th level (with respect to x). Denote the hitting time of the n-th level by

$$T_n = T_n(X_\cdot) := \inf\{t\ge 0 : (X_t - X_0)\cdot e_1 = n/\lambda_1\}, n\in\mathbb{Z}.$$

Also set

$$T_{\pm 0.5} := \inf\{t\ge 0 : (X_t - X_0)\cdot e_1 = \pm 0.5/\lambda_1\}.$$

Since $\ell_1 > 0$, the random walk is transient in the e_1 direction. Thus $(T_n)_{n\ge 0}$ are finite P_{ω^λ}-almost surely.

Proposition 4. *For any $n, m\in\mathbb{Z}^+$ and any balanced environment ω,*

$$P_{\omega^\lambda}(T_n < T_{-m}) = \frac{1-q_\lambda^m}{1-q_\lambda^{m+n}},$$

where $q_\lambda := (\frac{1-\lambda\ell_1}{1+\lambda\ell_1})^{1/\lambda_1}$.

Proof: Observe that the jumps of $(X_n \cdot e_1)_{n\geq 0}$ are lazy random walks on $\mathbb{Z}$, with the ratio of the probabilities of left-jump to right-jump equals $(1-\lambda\ell_1)/(1+\lambda\ell_1)$. Hence for $i,j \in \mathbb{Z}^+$,

$$P_{\omega^\lambda}(\tilde{T}_i < \tilde{T}_{-j}) = \frac{1-(\frac{1-\lambda\ell_1}{1+\lambda\ell_1})^j}{1-(\frac{1-\lambda\ell_1}{1+\lambda\ell_1})^{i+j}},$$

where $\tilde{T}_k := \inf\{n \geq 0 : (X_n - X_0)\cdot e_1 = k\}, k \in \mathbb{Z}$. The proposition follows by noting that $T_n = \tilde{T}_{n/\lambda_1}$. □

Lemma 5. *For all $\lambda \in (0,1), m \in \mathbb{N}$ and any balanced environment ω with ellipticity constant $\kappa \in (0, 1/(2d))$, if $t \geq 2m/\kappa$, then*

$$P_{\omega^\lambda}(T_m \geq t/\lambda_1^2) \leq e^{-t\kappa^2/8}.$$

Proof: Recall the definition of Y_n in (1). Note that

$$Y_n \cdot e_1 \geq 2\kappa\lambda_1 n - X_n \cdot e_1.$$

Since $Y_n \cdot e_1$ is a P_{ω^λ}-martingale with increments bounded by 2, we get

$$\begin{aligned}
&P_{\omega^\lambda}(T_m \geq t/\lambda_1^2)\\
&\leq P_{\omega^\lambda}(Y_{\lfloor t/\lambda_1^2\rfloor}\cdot e_1 \geq 2\kappa\lambda_1\lfloor t/\lambda_1^2\rfloor - m/\lambda_1)\\
&\leq P_{\omega^\lambda}(Y_{\lfloor t/\lambda_1^2\rfloor}\cdot e_1 \geq \kappa t/\lambda_1)\\
&\leq \exp\big(-\frac{(\kappa t/\lambda_1)^2}{8\lfloor t/\lambda_1^2\rfloor}\big) \leq \exp(-\frac{\kappa^2 t}{8}),
\end{aligned}$$

where we used Azuma-Hoeffding's inequality in the second to last inequality. □

Proposition 6. *There exists a constant $C_0 = C_0(\kappa, d) > 0$ such that for any balanced environment ω with ellipticity constant $\kappa > 0$,*

$$P_{\omega^\lambda}(\max_{0\leq s\leq T_1}|X_s| \geq C_0/\lambda_1) < 0.5.$$

Proof: Since

$$\max_{0\leq s\leq T_1}|X_s| \leq \max_{0\leq s\leq T_1}|Y_s| + \lambda T_1,$$

we have

$$\begin{aligned}
&P_{\omega^\lambda}(\max_{0\leq s\leq T_1}|X_s| \geq C/\lambda_1)\\
&\leq P_{\omega^\lambda}(T_1 > 1/\lambda_1^2) + P_{\omega^\lambda}(\max_{0\leq s\leq T_1}|Y_s| \geq C/\lambda_1 - \lambda/\lambda_1^2).
\end{aligned}$$

The Proposition follows by Lemma 5 and by Azuma-Hoeffding's inequality. □

3.2 Harnack inequality and its application

Let a be a nonnegative function on $\mathbb{Z}^d \times \mathbb{Z}^d$ such that for any x, $a(x,y) > 0$ only if x and y are neighbours, i.e., $|x-y| = 1$, denoted $x \sim y$. We also assume that

$$\sum_y a(x,y) = 1, \quad \forall x \in \mathbb{Z}^d.$$

Define the linear operator L_a acting on the set of functions on $\mathbb{Z}^d$ by

$$L_a f(x) = \sum_y a(x,y)(f(y) - f(x)).$$

Set

$$b(x) = \sum_y a(x,y)(y-x),$$

$$\text{and} \qquad b_0 = \sup |b|.$$

We assume that L_a is uniformly elliptic with constant $\kappa \in (0, \frac{1}{2d}]$. That is,

$$a(x,y) \geq \kappa \text{ for any } x, y \text{ such that } x \sim y.$$

For $r > 0, x \in \mathbb{R}^d$, let $B_r(x) = \{z \in \mathbb{Z}^d : |z - x| < r\}$. We also write $B_r(o)$ as B_r.

The following Harnack estimate is due to Kuo and Trudinger [9, Theorem 3.1]. See the Appendix for its proof.

Theorem 7 (Harnack inequality)**.** *Let u be a non-negative function on B_R, $R > 1$. If*

$$L_a u = 0$$

in B_R, then for any $\sigma \in (0,1)$ with $R(1-\sigma) > 1$, we have

$$\max_{B_{\sigma R}} u \leq C \min_{B_{\sigma R}} u,$$

where C is a positive constant depending on d, κ, σ and $b_0 R$.

With the Harnack inequality, we have

Lemma 8. *There exists a constant $c_1 \in (0,1]$ such that for any $\lambda \in (0,1)$, $x \in \mathbb{Z}^d$ and balanced environment ω,*

$$P^x_{\omega^\lambda}(X_{T_1} = \cdot) \geq c_1 P^{x+0.5e_1/\lambda_1}_{\omega^\lambda}(X_{T_{0.5}} = \cdot | T_{0.5} < T_{-0.5}). \tag{9}$$

Proof: For any $x \in \mathbb{Z}^d$, let

$$\mathcal{H}^x_{0.5} := \{y \in \mathbb{Z}^d : (y-x) \cdot e_1 = 0.5/\lambda_1\}.$$

Fix $w \in \mathcal{H}_1^x$. Then the function

$$f(z) := P^z_{\omega^\lambda}(X_\cdot \text{ visits } \mathcal{H}_1^x \text{ for the first time at } w)$$

satisfies

$$L_{\omega^\lambda} f(z) = 0$$

for all $z \in \{y : (y-x)\cdot e_1 < 1/\lambda_1\}$. By Theorem 7 (in this case $a = \omega^\lambda, R = 0.5/\lambda_1$ and $b_0 \le \lambda$), there exists a constant C_2 such that for any $y, z \in \mathcal{H}_{0.5}^x$ with $|z-y| < 0.5/\lambda_1$,

$$f(z) \ge C_2 f(y).$$

Hence for any $z \in \mathcal{H}_{0.5}^x$ such that $|z - (x + 0.5e_1/\lambda_1)| < C_0/\lambda_1$, we have

$$f(z) \ge C_2^{2C_0} f(x + 0.5e_1/\lambda_1). \tag{10}$$

Therefore,

$$\begin{aligned} P^x_{\omega^\lambda}(X_{T_1} = w) &\ge \sum_{|y-x|<C_0/\lambda} P^x_{\omega^\lambda}(X_{T_{0.5}} = y) P^y_{\omega^\lambda}(X_{T_{0.5}} = w) \\ &\overset{(10)}{\ge} C P^x_{\omega^\lambda}(|X_{T_{0.5}} - x| < C_0/\lambda_1) P^{x+0.5e_1/\lambda_1}_{\omega^\lambda}(X_{T_{0.5}} = w) \\ &\ge c_1 P^{x+0.5e_1/\lambda_1}_{\omega^\lambda}(X_{T_{0.5}} = w | T_{0.5} < T_{-0.5}) \end{aligned}$$

where in the last inequality we used the facts that (by Proposition 6)

$$P^x_{\omega^\lambda}(|X_{T_{0.5}} - x| < C_0/\lambda_1) > \frac{1}{2}$$

and

$$P^{x+0.5e_1/\lambda_1}_{\omega^\lambda}(T_{0.5} < T_{-0.5}) > \frac{1}{2}. \quad \square$$

3.3 Construction of the regeneration times

Let

$$\mu^x_{\omega^\lambda,1}(\cdot) = P^{x+0.5e_1/\lambda_1}_{\omega^\lambda}(X_{T_{0.5}} = \cdot | T_{0.5} < T_{-0.5}).$$

Recall that c_1 is the constant in Lemma 8. For any $\beta \in (0, c_1)$, we set

$$\mu^x_{\omega^\lambda,0}(\cdot) = \mu^{x,\beta}_{\omega^\lambda,0}(\cdot) := \left[P^x_{\omega^\lambda}(X_{T_1} = \cdot) - \beta \mu^x_{\omega^\lambda,1}(\cdot)\right]/(1-\beta).$$

Then by (9), both $\mu^x_{\omega^\lambda,1}$ and $\mu^x_{\omega^\lambda,0}$ are probability measures on $\mathcal{H}^x_{0.5}$ and

$$P^x_{\omega^\lambda}(X_{T_1} = u) = \beta \mu^x_{\omega^\lambda,1}(u) + (1-\beta)\mu^x_{\omega^\lambda,0}(u).$$

For any $\mathcal{O} \in \sigma(X_1, X_2, \ldots, X_{T_1}), x \in \mathbb{Z}^d$ and $i \in \{0,1\}$, put

$$\begin{aligned} \nu^x_{\omega^\lambda,i}(\mathcal{O}) &= \nu^{x,\beta}_{\omega^\lambda,i}(\mathcal{O}) \\ &:= \sum_y \left[i\mu^x_{\omega^\lambda,1}(y) + (1-i)\mu^x_{\omega^\lambda,0}(y)\right] P^x_{\omega^\lambda}(\mathcal{O} | X_{T_1} = y). \end{aligned} \tag{11}$$

Notice that under the environment measure P,

$$\nu^x_{\omega^\lambda,1}(X_{T_1} \in \cdot) = \mu^x_{\omega^\lambda,1}(\cdot)$$

is independent of $\sigma(\omega_y : y \cdot e_1 \le x \cdot e_1)$.

Now we will define the regeneration times.

We first sample a sequence $(\epsilon_i)_{i=1}^\infty \in \{0,1\}^{\mathbb{N}}$ of iid Bernoulli random variables according to the law Q_β defined by

$$Q_\beta(\epsilon_i = 1) = \beta \text{ and } Q_\beta(\epsilon_i = 0) = 1 - \beta.$$

Then, fixing $\epsilon := (\epsilon_i)_{i=1}^\infty$, we will define a new law $P_{\omega^\lambda,\epsilon}$ on the paths as follows (see Figure 1). For $x \in \mathbb{Z}^d$, set

$$P^x_{\omega^\lambda,\epsilon}(X_0 = x) = 1.$$

Assume that the $P^x_{\omega^\lambda,\epsilon}$-law for paths of length$\le n$ is defined. For any path $(x_i)_{i=0}^{n+1}$ with $x_0 = x$, define

$$\begin{aligned}&P^x_{\omega^\lambda,\epsilon}(X_{n+1} = x_{n+1}, \dots, X_0 = x_0)\\ &:= P^x_{\omega,\epsilon}(X_I = x_I, \dots, X_0 = x_0)\nu^{x_I}_{\omega^\lambda,\epsilon_J}(X_{n+1-I} = x_{n+1}, \dots, X_1 = x_{I+1}),\end{aligned}$$

where

$$J = J(x_0, \dots, x_n) := \max\{j \ge 0 : \mathcal{H}^{x_0}_j \cap \{x_i, 0 \le i \le n\} \ne \emptyset\}$$

is the highest level visited by $(x_i)_{i=0}^n$ and

$$I = I(x_0, \dots, x_n) := \min\{0 \le i \le n : x_i \in \mathcal{H}^{x_0}_J\}$$

is the hitting time to the J-th level. By induction, the law $P^x_{\omega^\lambda,\epsilon}$ is well-defined for paths of all lengths.


get 1
Go to the next level, following law $\nu_{\omega^\lambda,1}$
Flip a coin
get 0
Go to the next level, following law $\nu_{\omega^\lambda,0}$

Figure 1: The law $\bar{P}_{\omega^\lambda,\epsilon}$ for the walks.

Note that a path sampled by $P^x_{\omega^\lambda,\epsilon}$ is not a Markov chain, but the law of $X.$ under

$$\bar{P}^x_{\omega^\lambda} = \bar{P}^x_{\omega^\lambda,\beta} := Q_\beta \otimes P^x_{\omega^\lambda,\epsilon}$$

coincides with $P^x_{\omega^\lambda}$. That is,

$$\bar{P}^x_{\omega^\lambda}(X_\cdot \in \cdot) = P^x_{\omega^\lambda}(X_\cdot \in \cdot).$$

We denote by $\bar{\mathbb{P}}_\lambda = \bar{\mathbb{P}}_{\lambda,\beta} := P \otimes \bar{P}_{\omega^\lambda,\beta}$ the law of the triple $(\omega, \epsilon, X_\cdot)$. Expectations with respect to $\bar{P}^x_{\omega^\lambda}$ and $\bar{\mathbb{P}}_\lambda$ are denoted by $\bar{E}^x_{\omega^\lambda}$ and $\bar{\mathbb{E}}_\lambda (= \bar{\mathbb{E}}_{\lambda,\beta})$, respectively.

Next, for a path $(X_n)_{n\geq 0}$ sampled according to $P^o_{\omega^\lambda,\epsilon}$, we will define the regeneration times. See Figure 4 for an illustration.


Keep walking until reaching a level where the coin is "1"
Proceed to the next level (following $\nu_{\omega^\lambda,1}$) and ask:
will the path backtrack to the previous level in the future?
N
Y
Get a regeneration

Figure 2: The definition of a regeneration time.

To be specific, put $S_0 = 0, M_0 = 0$, and define inductively

$$\begin{aligned}
&S_{k+1} = \inf\{T_{n+1} : n/\lambda_1 \geq M_k \text{ and } \epsilon_n = 1\},\\
&R_{k+1} = S_{k+1} + T_{-1} \circ \theta_{S_{k+1}},\\
&M_{k+1} = X_{S_{k+1}} \cdot e_1 + N \circ \theta_{S_{k+1}}, \qquad k \geq 0.
\end{aligned}$$

Here θ_n denotes the time shift of the path, i.e, $\theta_n X_\cdot = (X_{n+i})_{i=0}^\infty$, and

$$N := \inf\{n/\lambda_1 : n/\lambda_1 > (X_i - X_0) \cdot e_1 \text{ for all } i \leq T_{-1}\}.$$

Set

$$\begin{aligned}
&&K := \inf\{k \geq 1 : S_k < \infty, R_k = \infty\},\\
&&\tau_1 := S_K,\\
&\text{and} &\tau_{k+1} = \tau_k + \tau_1 \circ \theta_{\tau_k}.
\end{aligned}$$

We call $(\tau_k)_{k\geq 1}$ the (β-)*regeneration times*. Intuitively, under $\bar{P}^x_{\omega^\lambda}$, whenever the walker visits a new level $\mathcal{H}_i, i \geq 0$, he flips a coin ϵ_i. If $\epsilon_i = 0$ (or 1), he then walks following the law $\nu_{\omega^\lambda,0}$ (or $\nu_{\omega^\lambda,1}$) until he reaches the $(i+1)$-th level. The regeneration time τ_1 is defined to be the first time of visiting a new level $\mathcal{H}_k$ such that the outcome ϵ_{k-1} of the previous coin-tossing is "1" and the path will never backtrack to the level $\mathcal{H}_{k-1}$ in the future. See Figure 3.



Figure 3: In this picture, $K = 2, X_{\tau_1} = 5/\lambda_1, M_1 = 4/\lambda_1$.

3.4 The renewal property of the regenerations

The regeneration times possess good renewal properties in the following sense:

1. Since the ratio of the probabilities of left-jump and right-jump of the lazy random walks $(X_n \cdot e_1)_{n\geq 0}$ (in $\mathbb{Z}$) is $(1-\lambda\ell_1)/(1+\lambda\ell_1)$, the law of $(X_{\tau_n} \cdot e_1)_{n\geq 1}$ does not depend on the environment ω. (Indeed, if we only observe the chain $(X_n \cdot e_1)_{n\geq 0}$ at the times when it *moves* and forget about its laziness, we get a random walk on $\mathbb{Z}$ with probabilities $(1-\lambda\ell_1)/2$ and $(1+\lambda\ell_1)/2$ of jumping to the left and to the right, respectively.) Furthermore, under $\bar{P}_{\omega^\lambda}$, the inter-regeneration distances $(e_1 \cdot X_{\tau_1} \circ \theta_{\tau_n})_{n=1}^{\infty}$ in the direction e_1 are iid random variables which are independent of $X_{\tau_1} \cdot e_1$, and
$$\bar{P}_{\omega^\lambda}(e_1 \cdot X_{\tau_1} \circ \theta_{\tau_n} \in \cdot) = \bar{P}_{\omega^\lambda}(X_{\tau_1} \cdot e_1 \in \cdot | T_{-1} = \infty), n \geq 1.$$

2. For $k \geq 0$, define
$$\begin{aligned} &\tilde{S}_{k+1} := \inf\{T_n : n/\lambda \geq M_k \text{ and } \epsilon_n = 1\},\\ &\tilde{\tau}_1 := \tilde{S}_K,\\ \text{and} \qquad &\tilde{\tau}_{k+1} := \tau_k + \tilde{\tau}_1 \circ \theta_{\tau_k}. \end{aligned}$$
Note that for $k \geq 1$,
$$\begin{aligned} &S_k = \tilde{S}_k + T_1 \circ \theta_{\tilde{S}_k}\\ \text{and} \qquad &X_{\tau_k} \cdot e_1 = X_{\tilde{\tau}_k} \cdot e_1 + 1/\lambda_1. \end{aligned}$$

Conditioning on $X_{\tilde{\tau}_k} = x$, the law of X_{τ_k} is $\mu^x_{\omega^\lambda,1}$, which is independent (under the environment measure P) of $\sigma(\omega_y : y \cdot e_1 \le x \cdot e_1)$. Moreover, after time τ_k, the path will never visit $\{y : y \cdot e_1 \le x \cdot e_1\}$. Thus the movement of the path after time τ_k is independent (under $\bar{\mathbb{P}}_\lambda$) of $(X_n)_{n \le \tilde{\tau}_k}$, and therefore, we expect

$$(\tilde{\tau}_1 \circ \theta_{\tau_k})_{k \ge 1}$$

to be iid random variables under $\bar{\mathbb{P}}_\lambda$. See Proposition 10 for a rigorous proof.

3. Although $(X_{\tau_1} \circ \theta_{\tau_k})_{k \ge 1}$ and $(\tilde{\tau}_1 \circ \theta_{\tau_k})_{k \ge 1}$ are both iid sequences, the inter-regeneration times $(\tau_1 \circ \theta_{\tau_k})_{k \ge 1}$ are not even independent. However, letting

$$\Delta_k := T_1 \circ \theta_{X_{\tilde{\tau}_k}} = \tau_k - \tilde{\tau}_k \text{ for } k \ge 1,$$

we can show that for every $k \ge 1$, $\lambda_1^2 \Delta_k$ is bounded by a constant plus an exponential random variable. So Δ_k is much less than $\tau_1 \circ \theta_{\tau_k}$, which is roughly $C/(\beta\lambda_1^2)$ (as will be shown in Proposition 14). In this sense, the inter-regeneration times $\tau_1 \circ \theta_{\tau_k}$ are *almost* iid if β is sufficiently small.

The rest of this subsection is devoted to the proof that $(\tilde{\tau}_1 \circ \theta_{\tau_k})_{k \ge 1}$ are iid (Proposition 10) and that Δ_k's are dominated by iid random variables of sizes $1/\lambda^2$ (Proposition 11).

We introduce the σ-field

$$\mathcal{G}_k := \sigma\big(\tilde{\tau}_k, (X_i)_{i \le \tilde{\tau}_k}, (\omega_y)_{y \cdot e_1 \le X_{\tilde{\tau}_k} \cdot e_1}\big).$$

Lemma 9. *For any appropriate measurable sets B_1, B_2 and any event*

$$B := \{(X_i)_{i \ge 0} \in B_1, (\omega_y)_{y \cdot e_1 > -1/\lambda_1} \in B_2\},$$

we have for $k \ge 1$,

$$\bar{\mathbb{P}}_\lambda(B \circ \bar{\theta}_{\tau_k} | \mathcal{G}_k) = \frac{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(B \cap \{T_{-1} = \infty\})\big]}{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(T_{-1} = \infty)\big]}.$$

Here $\bar{\theta}_n$ is the shift defined by

$$B \circ \bar{\theta}_n = \{(X_i)_{i \ge n} \in B_1, (\omega_y)_{(y - X_n) \cdot e_1 > -1/\lambda_1} \in B_2\}.$$

Proof: For simplicity, let us consider the case $k = 1$. We use θ^n to denote the shift of the ϵ-coins, i.e., $\theta^n \epsilon_\cdot = (\epsilon_i)_{i \geq n}$. For any $A \in \mathcal{G}_1$,

$$\begin{aligned}
&\bar{\mathbb{P}}_\lambda(B \circ \bar{\theta}_{\tau_1} \cap A) \\
&= E_{P\otimes Q_\beta}\big[\sum_{k\geq 1, x} P_{\omega^\lambda,\epsilon}(A \cap \{\tilde{S}_k < \infty, R_k = \infty, X_{\tilde{S}_k} = x\} \cap B \circ \bar{\theta}_{S_k})\big] \\
&= E_{P\otimes Q_\beta}\big[\sum_{k\geq 1, x, y} P_{\omega^\lambda,\epsilon}(A \cap \{\tilde{S}_k < \infty, X_{\tilde{S}_k} = x\}) \nu^x_{\omega^\lambda,1}(X_{T_1} = x + y) \\
&\qquad\qquad\qquad \times P^{x+y}_{\omega^\lambda, \theta^{k+1}\epsilon}(B \cap \{T_{-1} = \infty\})\big].
\end{aligned}$$

Note that in the last equality,

$$P_{\omega^\lambda,\epsilon}(A \cap \{\tilde{S}_k < \infty, X_{\tilde{S}_k} = x\})$$

is $\sigma\big((\epsilon_i)_{i\leq k}, (\omega_z)_{(z-x)\cdot e_1 \leq 0}\big)$- measurable, whereas

$$\nu^x_{\omega^\lambda,1}(X_{T_1} = x + y) P^{x+y}_{\omega^\lambda,\theta^{k+1}\epsilon}(B \cap \{T_{-1} = \infty\})$$

is $\sigma\big((\epsilon_i)_{i\geq k+1}, (\omega_z)_{(z-x)\cdot e_1 > 0}\big)$- measurable for $y \in \mathcal{H}^x_1$. Hence they are independent under $P \otimes Q_\beta$ and we have

$$\begin{aligned}
&\bar{\mathbb{P}}_\lambda(B \circ \bar{\theta}_{\tau_1} \cap A) \\
&= \sum_{k\geq 1} \bar{\mathbb{P}}_\lambda(A \cap \{\tilde{S}_k < \infty\}) E_P\big[\sum_y \nu_{\omega^\lambda,1}(X_{T_1} = y) \bar{P}^y_{\omega^\lambda}(B \cap \{T_{-1} = \infty\})\big].
\end{aligned} \tag{12}$$

Substituting B with the set of all events, we get

$$\bar{\mathbb{P}}_\lambda(A) = \sum_{k\geq 1} \bar{\mathbb{P}}_\lambda(A \cap \{\tilde{S}_k < \infty\}) E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(T_{-1} = \infty)\big]. \tag{13}$$

(12) and (13) yield that

$$\bar{\mathbb{P}}_\lambda(B \circ \bar{\theta}_{\tau_1} | A) = \frac{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(B \cap \{T_{-1} = \infty\})\big]}{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(T_{-1} = \infty)\big]}.$$

The lemma is proved for the case $k = 1$. The general case $k > 1$ follows by induction. □

The following proposition is an immediate consequence of Lemma 9.

Proposition 10. *Under* $\bar{\mathbb{P}}_\lambda$, $\tilde{\tau}_1, \tilde{\tau}_1 \circ \theta_{\tau_1}, \ldots, \tilde{\tau}_1 \circ \theta_{\tau_k}, \ldots$ *are independent random variables. Furthermore,* $(\tilde{\tau}_1 \circ \theta_{\tau_k})_{k\geq 1}$ *are iid with law*

$$\bar{\mathbb{P}}_\lambda(\tilde{\tau}_1 \circ \theta_{\tau_k} \in \cdot) = \frac{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(\tilde{\tau}_1 \in \cdot, T_{-1} = \infty)\big]}{E_P\big[\sum_y \mu_{\omega^\lambda,1}(y) \bar{P}^y_{\omega^\lambda}(T_{-1} = \infty)\big]}.$$

Note that although the inter-regeneration times $(\tau_1 \circ \theta_{\tau_k})_{k\geq 1}$ are not independent. However, the differences between $\tau_1 \circ \theta_{\tau_k}$ and $\tilde{\tau}_1 \circ \theta_{\tau_k}, k \geq 1$ are controlled by iid exponential random variables. Indeed, for any $x \in \mathbb{Z}^d, t \geq 2/\kappa$,

$$\begin{aligned}
&\nu^x_{\omega^\lambda,1}(\lambda_1^2 T_1 \geq t)\\
&\overset{(11)}{=} \sum_y \mu^x_{\omega^\lambda,1}(y) P^x_{\omega^\lambda}(\lambda_1^2 T_1 \geq t | X_{T_1} = y)\\
&\overset{(9)}{\leq} c_1^{-1} \sum_y P^x_{\omega^\lambda}(X_{T_1} = y) P^x_{\omega^\lambda}(\lambda_1^2 T_1 \geq t | X_{T_1} = y)\\
&= c_1^{-1} P^x_{\omega^\lambda}(\lambda_1^2 T_1 \geq t) \overset{\text{Lemma } 5}{\leq} c_1^{-1} e^{-t\kappa^2/8}.
\end{aligned}$$

Hence for $k \geq 1$,

$$P_{\omega^\lambda,\epsilon}(\lambda_1^2 \Delta_k \geq t | X_i, i \leq \tilde{\tau}_k) = \nu^{X_{\tilde{\tau}_k}}_{\omega^\lambda,1}(\lambda_1^2 T_1 \geq t) \leq c_1^{-1} e^{-t\kappa^2/8},$$

which implies that $\lambda_1^2 \Delta_k$ is stochastically dominated by an exponential random variable (with rate $\kappa^2/8$) plus a constant $c_2 := 2/\kappa - 8\kappa^{-2} \log c_1$. Thus we conclude:

Proposition 11. *Enlarging the probability space if necessary, one can couple $(\Delta_k)_{k\geq 1}$ with an iid sequence $(\xi_k)_{k\geq 1}$ such that each ξ_k is the sum of c_2 and an exponential random variable with rate $\kappa^2/8$, and that*

$$\lambda_1^2 \Delta_k \leq \xi_k, \text{ for all } k \geq 1.$$

Therefore, for any $n \geq 1$,

$$\tilde{\tau}_1 + \sum_{i=1}^{n-1} \tilde{\tau}_1 \circ \theta_{\tau_i} \leq \tau_n \leq \tilde{\tau}_1 + \sum_{i=1}^{n-1} \tilde{\tau}_1 \circ \theta_{\tau_i} + \sum_{i=1}^{n} \xi_i / \lambda_1^2. \tag{14}$$

4 Moment estimates

Set $\tau_0 = 0$. We will show that the typical values of $e_1 \cdot X_1 \circ \theta_{\tau_k}$ and $\tau \circ \theta_{\tau_k}$, $k \geq 0$ are $C/(\beta\lambda)$ and $C/(\beta\lambda^2)$, respectively.

Theorem 12. *Then there exists a constant $c_3 > 0$ such that for any elliptic balanced environment ω and any $\lambda, \beta \in (0, 1)$,*

$$\bar{E}_{\omega^\lambda} \exp(c_3 \beta \lambda_1 X_{\tau_1} \cdot e_1) \leq C.$$

Proof: For $0 \leq k \leq K - 1$, set

$$L_{k+1} = \inf\{n \geq \lambda_1 M_k : \epsilon_n = 1\} - \lambda_1 M_k + 1.$$

Then L_1 is the number of coins tossed to get the first '1' and

$$X_{S_1} \cdot e_1 = L_1/\lambda_1.$$

Moreover, for $1 \le k \le K-1$, let

$$N_k = N \circ \theta_{S_k}.$$

Then

$$(X_{S_{k+1}} - X_{S_k}) \cdot e_1 = N_k + L_{k+1}/\lambda_1, \quad k \ge 1.$$

So

$$X_{\mathcal{T}_1} \cdot e_1 = \sum_{i=1}^{K} L_i/\lambda_1 + \sum_{i=1}^{K-1} N_i. \tag{15}$$

First, we will compute the exponential moment of $L_i, i \le K$. Since $(L_i)_{i \ge 1}$ depends only on the coins $(\epsilon_i)_{i \ge 0}$, it is easily seen that they are iid geometric random variables with parameter β. Hence for any $s \in (0,1)$ (note that $(1-\beta)e^{s\beta} < e^{-(1-s)\beta} < 1$),

$$\bar{E}_{\omega^\lambda}[e^{s\beta L_i}] = \sum_{n=0}^{\infty} e^{s\beta(n+1)}(1-\beta)^n \beta = \frac{\beta e^{s\beta}}{1-(1-\beta)e^{s\beta}} \le \frac{1}{1-s}. \tag{16}$$

Next, we will compute the exponential moment of $N_i, i \le K-1$. By Proposition 4, putting

$$p_\lambda := \bar{P}_{\omega^\lambda}(T_{-1} = \infty) = 1 - q_\lambda,$$

we have

$$\begin{aligned}
&\bar{P}_{\omega^\lambda}(N = (n+1)/\lambda_1) \\
&= \bar{P}_{\omega^\lambda}(T_n < T_{-1} < T_{n+1}) \\
&= \bar{P}_{\omega^\lambda}(T_n < T_{-1}) - \bar{P}_{\omega^\lambda}(T_{n+1} < T_{-1}) \\
&= \frac{p_\lambda}{1-q_\lambda^{n+1}} - \frac{p_\lambda}{1-q_\lambda^{n+2}} \le q_\lambda^{n+1}, \quad n \ge 0.
\end{aligned}$$

Observe that conditioning on K, $(N_i)_{1 \le i < K}$ are iid under $\bar{P}_{\omega^\lambda}$. Hence

$$\bar{P}_{\omega^\lambda}(N_i = (n+1)/\lambda_1 | K > i) = \bar{P}_{\omega^\lambda}(N = (n+1)/\lambda_1 | T_{-1} < \infty) \le q_\lambda^n,$$

and for $s \in (0,1)$,

$$\bar{E}_{\omega^\lambda}[e^{s\beta\lambda_1 N_i} | K > i] \le \frac{e^{s\beta}}{1 - e^{s\beta} q_\lambda}. \tag{17}$$

Finally, note that under $\bar{P}_{\omega^\lambda} = Q_\beta \otimes P^o_{\omega^\lambda,\epsilon}$, K is a geometric random variable with success parameter p_λ, and $(L_i)_{1 \le i \le K}$ and $(N_i)_{1 \le i \le K}$ are iid sequences when conditioned on K. Therefore, by (15), (16) and (17), for $s \in (0,1)$,

$$\begin{aligned}\bar{E}_{\omega^\lambda} \exp(s\beta\lambda_1 X_{\tau_1} \cdot e_1) &\le \bar{E}_{\omega^\lambda}\left[\frac{1}{(1-s)^K}\left(\frac{e^{s\beta}}{1-e^{s\beta}q_\lambda}\right)\right] \\ &= \frac{p_\lambda}{1-s}\sum_{n=0}^{\infty}\left[\frac{e^{s\beta}q_\lambda}{(1-s)(1-e^{s\beta}q_\lambda)}\right]^n.\end{aligned}$$

The theorem follows by taking s small enough and noting that $q_\lambda < e^{-1}$. ☐

Corollary 13. *For P-almost every ω and any $\lambda, \beta \in (0,1)$,*

$$\bar{P}_{\omega^\lambda}(\beta\lambda_1^2\tau_1 \ge t) \le Ce^{-ct}.$$

Proof: By Lemma 5 and Theorem 12, for any $n \le \kappa t/(2\beta)$,

$$\begin{aligned}&\bar{P}_{\omega^\lambda}(\beta\lambda_1^2\tau_1 \ge t) \\ &\le \bar{P}_{\omega^\lambda}(X_{\tau_1} \cdot e_1 \ge n/\lambda_1) + \bar{P}_{\omega^\lambda}(\beta\lambda_1^2 T_n \ge t) \\ &\le Ce^{-c_3 n\beta} + e^{-\kappa^2 t/(8\beta)}.\end{aligned}$$

The Corollary follows by taking $n = \lfloor t/\beta \rfloor$. ☐

It follows from Corollary 13 and Lemma 9 (and noting that $P_{\omega^\lambda}(T_{-1} = \infty) = p_\lambda > 1/2$) that for $k \ge 1$,

$$\bar{\mathbb{P}}_\lambda(\beta\lambda_1^2\tau_1 \circ \theta_{\tau_k} \ge t) \le Ce^{-ct}. \tag{18}$$

Hence by Theorem 12, Corollary 13 and (18), we conclude that for any $p \ge 1, k \ge 0$, there exists a constant $C(p) < \infty$ such that

$$\bar{\mathbb{E}}_\lambda(\beta\lambda_1^2\tau_1 \circ \theta_{\tau_k})^p < C(p), \tag{19}$$

$$\text{and} \qquad \bar{\mathbb{E}}_\lambda(\beta\lambda_1 X_{\tau_1} \circ \theta_{\tau_k})^p < C(p). \tag{20}$$

Moreover, since $\bar{\mathbb{P}}_\lambda$-almost surely,

$$v_\lambda \cdot e_1 = \lim_{n\to\infty} \frac{X_{\tau_n} \cdot e_1}{\tau_n},$$

by (14) and the law of large numbers, we have

$$\frac{\bar{\mathbb{E}}_\lambda[e_1 \cdot X_{\tau_1} \circ \theta_{\tau_1}]}{\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1}] + E\xi_1/\lambda_1^2} \le v_\lambda \cdot e_1 \le \frac{\bar{\mathbb{E}}_\lambda[e_1 \cdot X_{\tau_1} \circ \theta_{\tau_1}]}{\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1}]}. \tag{21}$$

Proposition 14. *For any $\lambda, \beta \in (0,1)$,*

$$\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1}] \ge \frac{C}{\beta\lambda_1^2}. \tag{22}$$

Proof: By the definition of $L_i, i \geq 1$, we get

$$\bar{\mathbb{E}}_\lambda[e_1 \cdot X_{\tau_1} \circ \theta_{\tau_1}] \geq \bar{\mathbb{E}}_\lambda L_1/\lambda_1 \geq \frac{1}{\beta\lambda_1}. \tag{23}$$

On the other hand, by (2), $v_\lambda \cdot e_1 \leq \lambda$ for all $\lambda \in (0,1)$. This, together with (21) and (23), yields

$$\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1}] + E\xi_1/\lambda_1^2 \geq \frac{C}{\beta\lambda_1^2}.$$

Recalling (see Proposition 11) that $E\xi_1 \leq c_2 + 8/\kappa^2$, display (22) is proved for $\beta < (0, \beta_0)$, where β_0 is a sufficiently small constant.

It remains to prove (22) for $\beta \in [\beta_0, 1)$. Since $(|X_n|^2 - n)_{n\in\mathbb{N}}$ is a martingale, we get $\bar{\mathbb{E}}_\lambda \tau_1 \geq \bar{\mathbb{E}}_\lambda T_1 = \bar{\mathbb{E}}_\lambda[|X_{T_1}|^2] \geq 1/\lambda_1^2$. Our proof is complete. □

5 Proof of the Einstein relation

For $n \geq 1$, let $\alpha_n = \alpha_n(\beta, \lambda) := \bar{\mathbb{E}}_\lambda \tau_n$. Note that by (19) and (22),

$$\frac{Cn}{\beta\lambda^2} \leq \alpha_n \leq \frac{C(1)n}{\beta\lambda^2}. \tag{24}$$

Lemma 15. *Let f be a function that satisfies* (3). *Then*

$$\left| E_{Q_\lambda} f - \frac{1}{\bar{\mathbb{E}}_\lambda \tau_n} \bar{\mathbb{E}}_\lambda \left[\sum_{i=0}^{\tau_n} f(\bar{\omega}_i) \right] \right| \leq C N_f A_f (\beta + 1/n) \qquad \forall n \in \mathbb{N}, \lambda, \beta \in (0,1).$$

Proof: The Lemma is trivial when $n \leq N_f$, so we only consider $n > N_f$. Recall that $\tau_0 = 0$. For $k \geq 0$, set

$$Z_k = Z_k(f) = \sum_{i=\tau_k}^{\tau_{k+1}-1} f(\bar{\omega}_i),$$
$$\tilde{Z}_k = \tilde{Z}_k(f) = \sum_{i=\tau_k}^{\tilde{\tau}_{k+1}-1} f(\bar{\omega}_i).$$

Then, $(\tilde{Z}_{N_f+k})_{k\geq 0}$ is an N_f-dependent sequence with identical distribution under $\bar{\mathbb{P}}_\lambda$, and

$$\tilde{Z}_k \leq Z_k \leq \tilde{Z}_k + A_f \xi_k/\lambda^2, \quad \bar{\mathbb{P}}_\lambda\text{-a.s.} \tag{25}$$

Hence, for $n > N_f$,

$$\begin{aligned} \frac{(n-N_f)\bar{\mathbb{E}}_\lambda \tilde{Z}_{N_f}}{(n-1)\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1} + C/\lambda_1^2] + \bar{\mathbb{E}}_\lambda \tau_1} &\leq \frac{\bar{\mathbb{E}}_\lambda \sum_{k=0}^{n-1} Z_k}{\bar{\mathbb{E}}_\lambda \tau_n} \\ &\leq \frac{(n-N_f)\bar{\mathbb{E}}_\lambda[\tilde{Z}_{N_f} + CA_f/\lambda_1^2] + A_f \alpha_{N_f}}{(n-1)\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1 \circ \theta_{\tau_1}]}. \end{aligned} \tag{26}$$

On the other hand, since the $\mathbb{P}_\lambda$-law of $\bar{\omega}_n$ converges weakly to Q_λ, we have

$$Q_\lambda f = \lim_{n\to\infty} \bar{\mathbb{E}}_\lambda\big[\frac{1}{n}\sum_{i=0}^{n-1} f(\bar{\omega}_i)\big]. \tag{27}$$

By (25) and the law of large numbers for finite-dependent sequence,

$$\frac{\bar{\mathbb{E}}_\lambda \tilde{Z}_{N_f}}{\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1\circ\theta_{\tau_1}] + C/\lambda_1^2} \le \lim_{n\to\infty} \bar{\mathbb{E}}_\lambda\big[\frac{1}{\tau_n}\sum_{k=0}^{n-1} Z_k\big] \le \frac{\bar{\mathbb{E}}_\lambda \tilde{Z}_{N_f} + CA_f/\lambda_1^2}{\bar{\mathbb{E}}_\lambda[\tilde{\tau}_1\circ\theta_{\tau_1}]}. \tag{28}$$

The lemma follows by combining (26), (27) and (28) and using the moment bound (24). ☐

Lemma 16. *Let f be a function that satisfies* (3). *Then*

$$\frac{1}{\alpha_n}\bar{\mathbb{E}}_\lambda\left|\sum_{i=0}^{\tau_n} f(\bar{\omega}_i) - \sum_{i=0}^{\alpha_n} f(\bar{\omega}_i)\right| \le CA_f(\beta + \frac{1}{\sqrt{n}}) \quad \forall n\in\mathbb{N}, \lambda,\beta\in(0,1).$$

Proof: Noting that the left hand side is less than

$$\frac{A_f}{\alpha_n}\bar{\mathbb{E}}_\lambda|\tau_n - \alpha_n| \le \frac{A_f}{\alpha_n}\sqrt{\operatorname{Var}\tau_n},$$

it suffices to show

$$\operatorname{Var}\tau_n \le (C\beta^2 + C/n)\alpha_n^2. \tag{29}$$

Indeed, put

$$A_n := \tilde{\tau}_1 + \sum_{i=1}^{n-1} \tilde{\tau}_1\circ\theta_{\tau_i}$$

and $B_n := A_n + \sum_{i=1}^n \xi_i/\lambda_1^2$. Then by (14), we have $A_n \le \tau_n \le B_n$. Thus

$$A_n - \bar{\mathbb{E}}_\lambda A_n - Cn/\lambda^2 \le \tau_n - \alpha_n \le B_n - \bar{\mathbb{E}}_\lambda B_n + Cn/\lambda^2.$$

Moreover,

$$A_n - \bar{\mathbb{E}}_\lambda A_n - C\beta\alpha_n \le \tau_n - \alpha_n \le B_n - \bar{\mathbb{E}}_\lambda B_n + C\beta\alpha_n,$$

and so

$$\begin{aligned}\operatorname{Var}\tau_n &\le \bar{\mathbb{E}}_\lambda(A_n - \bar{\mathbb{E}}_\lambda A_n - C\beta\alpha_n)^2 + \bar{\mathbb{E}}_\lambda(B_n - \bar{\mathbb{E}}_\lambda B_n + C\beta\alpha_n)^2\\ &= \operatorname{Var}A_n + \operatorname{Var}B_n + C\beta^2\alpha_n^2.\end{aligned}$$

Since (recall Proposition 10)

$$\operatorname{Var}A_n = \operatorname{Var}\tilde{\tau}_1 + (n-1)\operatorname{Var}\tilde{\tau}_1\circ\theta_{\tau_1} \overset{(19)}{\le} Cn(\beta\lambda^2)^{-2}$$

and

$$\operatorname{Var} B_n \le 2\big(\operatorname{Var} A_n + \operatorname{Var}(\sum_{i=1}^{n} \xi_i/\lambda_1^2)\big) = 2\operatorname{Var} A_n + Cn/\lambda_1^4 \le Cn(\beta\lambda^2)^{-2},$$

we conclude that

$$\operatorname{Var} \tau_n \le Cn(\beta\lambda^2)^{-2} + C\beta\alpha_n^2.$$

This completes the proof of (29). □

Proof of Theorem 3:
Since the left side is uniformly bounded (by $2A_f$) for all t, it suffices to prove the inequality for all sufficiently large t. Let

$$\beta = \beta(t) = t^{-1/3} \tag{30}$$

and set $n = n(t,\lambda)$ to be the integer that satisfies

$$\alpha_n \le \frac{t}{\lambda^2} < \alpha_{n+1}.$$

By (24), the existence of $n(t,\lambda)$ is guaranteed (since t is large). Moreover,

$$n \ge Ct\beta = Ct^{2/3}. \tag{31}$$

Since

$$\begin{aligned}
&\Big|\frac{\lambda^2}{t}\bar{\mathbb{E}}_\lambda \sum_{i=0}^{t/\lambda^2} f(\bar\omega_i) - \frac{1}{\alpha_n}\bar{\mathbb{E}}_\lambda \sum_{i=0}^{\alpha_n} f(\bar\omega_i)\Big| \\
&\le \Big|\frac{\lambda^2}{t}\bar{\mathbb{E}}_\lambda\big[\sum_{i=0}^{t/\lambda^2} f(\bar\omega_i) - \sum_{i=0}^{\alpha_n} f(\bar\omega_i)\big]\Big| + \Big|(\frac{\lambda^2}{t} - \frac{1}{\alpha_n})\bar{\mathbb{E}}_\lambda \sum_{i=0}^{\alpha_n} f(\bar\omega_i)\Big| \\
&\le A_f\bigg(\frac{\lambda^2}{t}\bar{\mathbb{E}}_\lambda[\alpha_{n+1} - \alpha_n] + (\frac{1}{\alpha_n} - \frac{1}{\alpha_{n+1}})\alpha_n\bigg) \\
&\le CA_f\bigg(\frac{1}{t\beta} + \frac{1}{n}\bigg),
\end{aligned}$$

combining Lemma 15, Lemma 16 and the above inequality, we conclude that if t is sufficiently large, then

$$\left|Q_\lambda f - \frac{\lambda^2}{t}\bar{\mathbb{E}}_\lambda \sum_{i=0}^{t/\lambda^2} f(\bar\omega_i)\right| \le CN_fA_f(\frac{1}{\sqrt{t\beta}} + \frac{1}{n} + \beta) \le \frac{CN_fA_f}{t^{1/3}}.$$

□

6 Discussions

1. Our arguments also work for non-nearest neighbour walks (with bounded size of jumps) in finite-dependent balanced environment.
2. Is the ER also true for mixing environments?
3. If we drop the uniform ellipticity assumption and assume only ellipticity, is the ER still true? Further, is it true for non-elliptic 'genuinely d-dimensional' balanced environment (see [2] for the definition)?
4. Is v_λ a monotone function of λ in an appropriate sense?
5. If the perturbation of the environment is not proportional to the original environment, what is $\mathrm{d}v_\lambda/\mathrm{d}\lambda$?
6. If the original environment is ballistic, what do we expect?
7. Can we get a Harnack inequality when the environment is non-elliptic 'genuinely d-dimensional'?

7 Acknowledgment

I thank my PhD advisor, Ofer Zeitouni, for introducing this problem to me and for numerous useful discussions. I thank Pierre Mathieu for many interesting discussions about the Einstein relation.

8 Appendix: Proof of the Harnack inequality

The purpose of this chapter is to present the proof the Harnack inequality (Theorem 7) for discrete harmonic functions. The Harnack inequality was used in Section 3 to construct the regeneration times. These inequalities are due to Kuo and Trudinger [9]. For the purpose of self-containedness, we will give the complete proofs of these estimates. We follow the arguments in [9], adding to it some extra details.

Recall the definitions of a, L_a, b, b_0 and the ellipticity assumption in Section 3.2.

8.1 Maximum principle

For any bounded set $E \subset \mathbb{Z}^d$, let $\partial E := \{y \in E^c : x \sim y \text{ for some } x \in E\}$, $\bar{E} = E \bigcup \partial E$ and $\mathrm{diam}(E) = \max\{|x-y|_\infty : x, y \in E\}$.

Theorem 17 (Maximum principle). *Let $E \subset \mathbb{Z}^d$ be bounded and u be a function on $\bar{E}$. For $x \in E$, define*

$$I_u(x) = \{s \in \mathbb{R}^d : u(x) - s \cdot x \geq u(z) - s \cdot z, \forall z \in \bar{E}\}.$$

If

$$L_a u(x) \geq -g(x)$$

for all $x \in E$ *such that* $I_u(x) = I_u(x, E, a) \neq \emptyset$, *then*

$$\max_E u \leq C \operatorname{diam}(\bar{E}) \big(\sum_{x \in E, I_u(x) \neq \emptyset} |g|^d \big)^{1/d} + \max_{\partial E} u,$$

where C *is a constant determined by* d, κ *and* $b_0 \operatorname{diam}(E)$.

Proof: Without loss of generality, we assume $g \geq 0$ and

$$\max_E u = u(x_0) > \max_{\partial E} u$$

for some $x_0 \in E$. Otherwise, there is nothing to prove.

For $s \in \mathbb{R}^d$ such that

$$\begin{aligned} |s|_\infty &\leq [u(x_0) - \max_{\partial E} u]/(d \operatorname{diam} \bar{E}) \\ &=: R = R(u, E), \end{aligned} \tag{32}$$

we have

$$u(x_0) - u(x) \geq s \cdot (x_0 - x)$$

for all $x \in \partial E$, which implies that $\max_{z \in \bar{E}} u(z) - s \cdot z$ is achieved in E. Hence $s \in \bigcup_{x \in E} I_u(x)$ and the cube

$$Q_R := \{x : |x|_\infty < R\} \subset \bigcup_{x \in E} I_u(x).$$

For any $p \in \mathbb{R}^d$, set

$$f(p) = (|p|^{d/d-1} + \mu^{d/d-1})^{1-d},$$

where $\mu > 0$ is a constant to be fixed later. Noting that for any $x \in E$, $I_u(x) \subset \mathbb{R}^d$ is bounded and closed, we can choose $p_x \in I_u(x)$ so that

$$|p_x| = \min_{p \in I_u(x)} |p|.$$

Then

$$f(p_x) = \max_{p \in I_u(x)} f(p).$$

Thus

$$\int_{Q_R} f(s) \, \mathrm{d}s \leq \int_{\bigcup_{x \in E} I_u(x)} f(s) \, \mathrm{d}s \leq \sum_{x : I_u(x) \neq \emptyset} f(p_x) |I_u(x)|, \tag{33}$$

where $|I_u(x)|$ denotes the Lebesgue measure of $I_u(x)$.

Further, we will show that for any $x \in E$ with $I_u(x) \neq \emptyset$,

$$|I_u(x)| \leq (2/\kappa)^d [g(x) + b(x)p_x]^d. \tag{34}$$

To this end, we fix an $x \in E$ with $I_u(x) \neq \emptyset$ and set

$$w(z) = u(z) - p_x(z - x), \quad \forall z \in \bar{E}.$$

Then $w(x) \geq w(z)$ for all $z \in \bar{E}$ and

$$I_u(x) = I_w(x) + p_x. \tag{35}$$

Since for any $q \in I_w(x)$ and $i = 1, \ldots, d$,

$$w(x) - w(x \pm e_i) \geq \mp q_i,$$

we obtain (by the ellipticity and by $w(x) \geq w(z)$, $\forall z \in \bar{E}$)

$$\begin{aligned} 0 \leq \kappa |q|_\infty &\leq \sum_y a(x,y)(w(x) - w(y)) \\ &= -L_a u + b(x)p_x \\ &\leq g(x) + b(x)p_x. \end{aligned}$$

Hence

$$I_w(x) \subset \big[\frac{-g(x) - b(x)p_x}{\kappa}, \frac{g(x) + b(x)p_x}{\kappa}\big]^d$$

and

$$|I_u(x)| \overset{(35)}{=} |I_w(x)| \leq (2/\kappa)^d [g(x) + b(x)p_x]^d.$$

(34) is proved.

(34) and (33) yield

$$\int_{Q_R} f(s)\,\mathrm{d}s \leq (\frac{2}{\kappa})^d \sum_{x : I_u(x) \neq \emptyset} f(p_x)[g(x) + b(x)p_x]^d.$$

Since by Hölder's inequality,

$$g(x) + |b(x)||p_x| \leq \big[(\frac{g(x)}{\mu})^d + |b(x)|^d\big]^{1/d} \big[\mu^{d/d-1} + |p_x|^{d/d-1}\big]^{(d-1)/d},$$

we get

$$\int_{Q_R} f(s)\,\mathrm{d}s \leq (\frac{2}{\kappa})^d \sum_{x : I_u(x) \neq \emptyset} \big[(\frac{g(x)}{\mu})^d + |b(x)|^d\big]. \tag{36}$$

On the other hand, by Hölder's inequality

$$f(s) = (|s|^{d/d-1} + \mu^{d/d-1})^{1-d} \geq 2^{2-d}(|s|^d + \mu^d)^{-1}.$$

Thus

$$\int_{Q_R} f(s)\,\mathrm{d}s \geq \int_{B_R} f(s)\,\mathrm{d}s \geq 2^{2-d}\int_{B_R} (|s|^d+\mu^d)^{-1}\,\mathrm{d}s$$
$$= 2^{2-d}\frac{\mathcal{O}_d}{d}\log[(\frac{R}{\mu})^d+1], \tag{37}$$

where $\mathcal{O}_d$ is the area of the unit sphere in $\mathbb{R}^d$.

Finally, combining (36) and (37) and putting

$$\mu := [\sum_{x:I_u(x)\neq\emptyset} g(x)^d]^{1/d},$$

we conclude that

$$\kappa^d 2^{2-2d}\frac{\mathcal{O}_d}{d}\log[(\frac{R}{\mu})^d+1] \leq 1+(b_0\operatorname{diam}\bar{E})^d.$$

Recalling the definition of $R = R(u, E)$ in (32), the theorem follows. □

8.2 Mean-value inequality

Let $u^+ = u \vee 0$. For any finite subset $E \subset \mathbb{Z}^d, p > 0$ and any function f on E, set

$$\|f\|_{E,p} := \big(|E|^{-1}\sum_{x\in E}|f(x)|^p\big)^{1/p}.$$

Lemma 18. *For fixed $R > 0$ and $\beta \geq 2$, let $\eta(x) = \eta_R(x) := (1 - |x|^2/R^2)^\beta 1_{|x|<R}$ be a function on $\mathbb{R}^d$. For any $\beta \geq 2$ and any function u on $\bar{B}_R$ such that*

$$L_a u \geq 0$$

in B_R, we let $v = \eta u^+$. Then for any $x \in B_R$ with $I_v(x) = I_v(x, B_R, a) \neq \emptyset$,

$$-L_a v(x) \leq C\eta^{1-2/\beta}R^{-2}u^+,$$

where the constant C depends only on β and b_0R.

Proof: We only need to consider the nontrivial case that $v \not\equiv 0$. For $s = s(x) \in I_v(x) \neq \emptyset$, recalling the definition of I_v one has

$$|s| \leq 2v(x)/(R-|x|).$$

Note that $I_v(x) \neq \emptyset$ implies $u(x) > 0$. If further $R^2 - |x|^2 \geq 4R$, computations as in [7] [1] reveal that for $y \sim x$,

$$2^{-\beta} \leq \frac{\eta(y)}{\eta(x)} \leq 2^\beta, \tag{38}$$

$$|\eta(x)-\eta(y)| \leq \beta 2^\beta R^{-1}\eta(x)^{1-1/\beta}, \tag{39}$$

$$|\eta(x)-\eta(y)-\nabla\eta(x)(x-y)| \leq \beta(\beta-1)2^\beta R^{-2}\eta(x)^{1-2/\beta}, \tag{40}$$

$$|s| \leq 4\eta^{1-1/\beta}R^{-1}u, \tag{41}$$

---

[1] Note that R^{-2} in [7, (26)] should be R^{-1}. This typo is corrected here, see (41).

where

$$\nabla\eta = -2\beta x R^{-2}\eta^{1-1/\beta} \tag{42}$$

is the gradient of η. For any $z \in \bar{E}$, set

$$w(z) = v(z) - s\cdot(z-x).$$

By the definition of s, we have $w(x) \geq w(z)$ for all $z \in \bar{E}$. Then

$$\begin{aligned} v(x)-v(y) = \frac{\eta(x)}{\eta(y)}(v(x)-v(y)) &+ \frac{\eta(y)-\eta(x)}{\eta(y)}s(x-y) \\ &+ \frac{\eta(y)-\eta(x)}{\eta(y)}(w(x)-w(y)). \end{aligned} \tag{43}$$

We first consider the case when $I_v(x) \neq \emptyset$ and $R^2-|x|^2 \geq 4R$. By (39), for any $y \sim x$,

$$\frac{\eta(y)-\eta(x)}{\eta(y)}(w(x)-w(y)) \leq \beta 2^{\beta} R^{-1}\eta(x)^{-1/\beta}\frac{\eta(x)}{\eta(y)}(w(x)-w(y)). \tag{44}$$

Since

$$\begin{aligned} &\sum_y a(x,y)\frac{\eta(x)}{\eta(y)}(w(x)-w(y)) \\ &= \sum_y a(x,y)\Big[\frac{\eta(x)}{\eta(y)}\big(v(x)-v(y)\big) + \frac{\eta(x)-\eta(y)}{\eta(y)}s(y-x)\Big] + s\cdot b(x), \end{aligned}$$

by (43) and (44), we obtain

$$\begin{aligned} &\sum_y a(x,y)(v(x)-v(y)) \\ &\leq (1+\beta 2^{\beta}R^{-1}\eta(x)^{-1/\beta})\sum_y a(x,y)\Big[\frac{\eta(x)}{\eta(y)}\big(v(x)-v(y)\big) + \frac{\eta(x)-\eta(y)}{\eta(y)}s(y-x)\Big] \\ &\quad + \beta 2^{\beta}R^{-1}\eta(x)^{-1/\beta}|s|b_0. \end{aligned} \tag{45}$$

Moreover, recalling that $u(x) > 0$ (because $I_v(x) \neq \emptyset$),

$$\begin{aligned} &\sum_y a(x,y)\frac{\eta(x)}{\eta(y)}\big(v(x)-v(y)\big) \\ &= \sum_y a(x,y)\left[\eta(x)\big(u(x)-u^+(y)\big) + \big(\eta(x)-\eta(y)\big)u(x) + \frac{(\eta(x)-\eta(y))^2}{\eta(y)}u(x)\right] \\ &\overset{a\geq 0}{\leq} -\eta(x)L_a u(x) + \sum_y a(x,y)\left[\big(\eta(x)-\eta(y)\big)u(x) + \frac{(\eta(x)-\eta(y))^2}{\eta(y)}u(x)\right] \\ &\overset{L_a u\geq 0}{\leq} \sum_y a(x,y)\left[\big(\eta(x)-\eta(y)-\nabla\eta(x)(x-y)\big)u(x) + \frac{(\eta(x)-\eta(y))^2}{\eta(y)}u(x)\right] \\ &\qquad - \nabla\eta(x)b(x)u(x) \\ &\leq \beta^2 2^{3\beta+1}\eta^{1-2/\beta}R^{-2}u + 2(b_0R)\beta\eta^{1-2/\beta}R^{-2}u, \end{aligned} \tag{46}$$

where we used (38), (39), (40) and (42) in the last inequality. Hence, by (45), (46) and noting $R^{-1}\eta^{-1/\beta} \le 1/4$, we conclude that

$$\begin{aligned} -L_a v &\le \beta^2 2^\beta (2^{3\beta} + b_0 R)\eta^{1-2/\beta} R^{-2} u + \beta 2^\beta R^{-1} \eta^{-1/\beta} |s| b_0 \\ &\overset{(41)}{\le} \beta^2 2^\beta (2^{3\beta} + 3 b_0 R)\eta^{1-2/\beta} R^{-2} u \end{aligned}$$

holds in $\{x : R^2 - |x|^2 \ge 4R, I_v(x) \ne \emptyset\}$.

On the other hand, if $R^2 - |x|^2 < 4R$, then $\eta^{1/\beta} \le 4/R$. Thus by the fact that $v \ge 0$, we have $-L_a v \le v(x) \le 16\eta^{1-2/\beta} R^{-2} u$. ☐

By the same argument as in the proof of [7, Theorem 12], Theorem 17 and Lemma 18 imply

Theorem 19 (Mean-value inequality)**.** *For any function u on $\bar{B}_R$ such that*

$$L_a u \ge 0, \quad x \in B_R$$

and any $\sigma \in (0,1)$, $0 < p \le d$, we have

$$\max_{B_{\sigma R}} u \le C \|u^+\|_{B_R, p},$$

where C depends on σ, p, κ, d and $b_0 R$.

8.3 Proof of the Harnack inequality

Lemma 20. *Suppose u is a non-negative function on $\bar{B}_R$ that satisfies*

$$L_a u \le 0$$

in B_R. Then for any $\sigma \le \tau < 1$,

$$\min_{B_{\tau R}} u \ge C \min_{B_{\sigma R}} u, \tag{47}$$

where C depends on κ, d, σ, τ and $b_0 R$.

Proof: Recall the definition of $\eta = \eta_R(x)$ in Lemma 18. We will first show that there exists a constant $\beta = \beta(\sigma, b_0 R, \kappa)$ such that

$$L_a \eta \ge -(2^\beta + \beta^3) R^{-3} \qquad \text{in } B_R \setminus B_{\sigma R}. \tag{48}$$

If $R - 1 \le |x| < R$, $\eta(x) \le (2/R)^\beta \le 2^\beta R^{-3}$ for $\beta \ge 3$. Hence for $\beta \ge 3$,

$$L_a \eta \ge -\eta \ge -2^\beta R^{-3}.$$

If $\sigma R \le |x| < R - 1$, then $y \in B_R$ for all $y \sim x$. Since for $i = 1, \ldots, d$, the third derivative $D_i^3 \eta$ of η with respect to x_i satisfies

$$\begin{aligned} |D_i^3 \eta| &= \left|4\beta(\beta-1) x_i R^{-4} \eta^{1-3/\beta} [3(1 - |x|^2/R^2) - 2(\beta-2) x_i^2/R^2]\right| \\ &\le 4\beta(\beta-1)(2\beta-1) R^{-3}, \end{aligned}$$

by Taylor's expansion, we have

$$\eta(x+e)-\eta(x)\ge \nabla\eta(x)\cdot e+\frac{1}{2}e^TD^2\eta(x)e-\frac{8}{6}\beta^3R^{-3}.$$

Thus

$$\begin{aligned}L_a\eta(x)&=\sum_e a(x,e)(\eta(x+e)-\eta(x))\\&\ge \nabla\eta\cdot b(x)+\frac{1}{2}\sum_e a(x,x+e)e^TD^2\eta(x)e-\frac{4}{3}\beta^3R^{-3}.\end{aligned}$$

Noting that

$$\nabla\eta\cdot b(x)\overset{(42),\eta\le 1}{\ge}-2(b_0R)\beta R^{-2}\eta^{1-2/\beta},$$

and for $\sigma R\le |x|<R-1$,

$$\begin{aligned}&\sum_e a(x,x+e)e^TD^2\eta(x)e\\&=\sum_{i=1}^d(a(x,x+e_i)+a(x,x-e_i))D_{ii}\eta(x)\\&=2\beta R^{-2}\eta^{1-2/\beta}\sum_{i=1}^d\big(a(x,x+e_i)+a(x,x-e_i)\big)\big(\frac{2(\beta-1)x_i^2}{R^2}-(1-\frac{|x|^2}{R^2})\big)\\&\ge 2\beta R^{-2}\eta^{1-2/\beta}[4\kappa(\beta-1)\sigma^2-1],\end{aligned}$$

we have

$$L_a\eta\ge[4\kappa(\beta-1)\sigma^2-1-2b_0R]R^{-2}\eta^{1-2/\beta}-\frac{4}{3}\beta^3R^{-3}.$$

Hence (48) also holds for $\sigma R\le|x|<R-1$ as we take

$$\beta\ge 1+\frac{1+2b_0R}{4\kappa\sigma^2}.$$

(48) is proved.

Next, let $m_\sigma:=\min_{B_{\sigma R}}u$ and $w:=m_\sigma\eta-u$. Then

$$\max_{B_{\tau R}}w\ge(1-\tau^2)^\beta m_\sigma-m_\tau. \tag{49}$$

Since $w\le 0$ in $B_{\sigma R}\bigcup B_R^c$, and

$$L_aw\overset{(48)}{\ge}-(2^\beta+\beta^3)m_\sigma R^{-3}\qquad\text{in } B_R/B_{\sigma R},$$

we get by the maximum principle that

$$\max_{B_R}w\le C_1m_\sigma R^{-1}, \tag{50}$$

where C_1 depends on κ, d, σ and b_0R. By (49) and (50),

$$[(1-\tau^2)^\beta - \frac{C_1}{R}]m_\sigma \le m_\tau.$$

Therefore, (47) holds if R satisfies

$$R > \frac{2C_1}{(1-\tau^2)^\beta}.$$

For $R \le \frac{2C_1}{(1-\tau^2)^\beta}$, it follows by iteration (note $\kappa u(x) \le u(y)$ for $x \sim y$) that

$$\kappa^{2C_1(1-\tau^2)^{-\beta}} m_\sigma \le m_\tau.$$

(47) is proved. ☐

For any $z \in \mathbb{Z}^d$ and any $n = (n_1, \ldots, n_d) \in \mathbb{N}^d$, we let

$$N(z,n) := (z + \prod_{i=1}^d [0, n_i - 1]) \cap \mathbb{Z}^d.$$

We say that $N(z,n)$ is *nice* if n satisfies $\max_{i,j}|n_i - n_j| \le 1$. Call $|n|_\infty$ the *length* of the nice rectangle $N(z,n)$. Intuitively, a nice rectangle is "nearly a cube".

Proposition 21. *Let u be a nonnegative function on $\bar{B}_R, R > 0$ such that*

$$L_a u \le 0 \quad \text{in } B_R.$$

Suppose $r \in (0, R/7\sqrt{d}]$ and $N = N(z,n) \subset Q_r$ is a nice rectangle in Q_r. Then there exists a constant $\delta = \delta(d, \kappa, b_0R) \in (0,1)$ such that if $\Gamma \subset B_R$ satisfies

$$|\Gamma \cap N| \ge \delta |N|,$$

then

$$\min_{N'} u \ge C \min_{\Gamma} u,$$

where $N' = (z + \prod_{i=1}^d [-n_i, 2n_i - 1]) \cap \mathbb{Z}^d$ and C depends on κ, d, σ, τ and b_0R.

Proof: When $|n|_\infty = 1$, N is a singleton, and the proposition follows by iteration (note that $u(x) \le \kappa u(y)$ for any $x \sim y$). So we only consider the case when the length of N is ≥ 2.

Denote the center of N by $O_N = z + (\frac{n_1-1}{2}, \cdots, \frac{n_d-1}{2}) \in (\frac{1}{2}\mathbb{Z})^d$. Setting $h =: \min_i n_i/2$, we have

$$B_h(O_N) \subset N \subset B_{2\sqrt{d}h}(O_N).$$



Figure 4: N is the rectangle in the center. $B_h(O_N)$ is the small circle.

Since $h \geq \frac{|n|_\infty - 1}{2} \geq \frac{|n|_\infty}{4}$, we have

$$N' \subset B_{3|n|_\infty \sqrt{d}/2}(O_N) \subset B_{6\sqrt{d}h}(O_N).$$

Suppose for some $\delta \in (0,1)$,

$$|\Gamma \cap N| \geq \delta |N|.$$

Let $u_\Gamma =: \min_\Gamma u$ and

$$v =: u_\Gamma - u,$$

then $L_a v \leq 0$ and $v^+|_\Gamma = 0$. By Theorem 19,

$$\begin{aligned}
\max_{B_{h/2}(O_N)} v &\leq C \frac{1}{|B_h|} \sum_{B_h(O_N)} v^+ \\
&\leq C \frac{|B_h(O_N) \setminus \Gamma|}{|B_h|} \max_{B_h(O_N)} v \\
&\overset{|B_h| \geq C|N|}{\leq} C \frac{|N \setminus \Gamma|}{|N|} \max_{B_h(O_N)} v \leq C_2(1-\delta) \max_{B_h(O_N)} v,
\end{aligned}$$

where C_2 depends on κ, d and $b_0 R$. Taking $\delta = \delta(\kappa, d, b_0 R)$ big enough such that $C_2(1-\delta) \leq 1/2$, we get

$$\max_{B_{h/2}(O_N)} v \leq \frac{1}{2} \max_{B_h(O_N)} v.$$

Hence

$$u_\Gamma - \min_{B_{h/2}(O_N)} u \le \frac{1}{2}(u_\Gamma - \min_{B_h(O_N)} u).$$

Therefore, noting that (since $r \le R/7\sqrt{d}$) $B_{7\sqrt{d}h}(O_N) \subset B_R$,

$$u_\Gamma \le 2 \min_{B_{h/2}(O_N)} u \overset{\text{Lemma 20}}{\le} C \min_{B_{6\sqrt{d}h}(O_N)} u \le C \min_{N'} u,$$

with C depends on κ, d and b_0R. □

Lemma 22. *Let u be a nonnegative function on $\bar{B}_R, R > 0$ such that*

$$L_a u \le 0 \quad \text{in } B_R.$$

Let $r \in (0, R/7\sqrt{d}]$. Then for any $\Gamma \subset Q_r$, there exists a subset $\Gamma_\delta \supset \Gamma$ of Q_r such that either $\Gamma_\delta = Q_r$ or $|\Gamma_\delta| > \delta^{-1}|\Gamma|$ holds, and

$$\min_{\Gamma_\delta} u \ge \gamma \min_{\Gamma} u.$$

Here the constant γ depends only on κ, d and b_0R, and δ is the same as in Proposition 21.

Proof: We will construct Γ_δ through a cube decomposition procedure.

Observe that any nice rectangle with length $l \ge 2$ can be decomposed into (at most 2^d) smaller disjoint nice rectangles whose lengths are either $\lfloor \frac{l}{2} \rfloor$ or $\lfloor \frac{l}{2} \rfloor + 1$. With abuse of terminology, we say that such a decomposition is *nice*. Note that a nice decomposition may not be unique.

For any $\Gamma \subset Q_r$, set

$$\mathcal{N} = \mathcal{N}(\Gamma) := \{N : N \text{ is nice and } |\Gamma \cap N| \ge \delta |N|\}.$$

Now perform cube decompositions to Q_r as follows. Assume that we have an imaginary "bag". In the first step, we put Q_r into our "bag" if $Q_r \in \mathcal{N}$, and decompose Q_r nicely (into at most 2^d nice rectangles) if otherwise. In the second step, we we repeat the same procedure on each of the remaining rectangles, i.e. put a rectangle into our "bag" if it is in $\mathcal{N}$, and decompose a rectangle (of lengths≥ 2) nicely if it is not in $\mathcal{N}$. So on and so forth. We will stop if we have nothing to decompose or all the remaining rectangles are singletons in $Q_r \setminus \Gamma$. The process will end within finite steps. Denote the collection of the rectangles in our "bag" by $\mathcal{N}_0 (\subset \mathcal{N})$.

For $N \in \mathcal{N}_0$ and $N \ne Q_r$, we denote by N^{-1} its *prior*, i.e, N is obtained from a nice decomposition of N^{-1} in the previous step. Set $Q_r^{-1} = Q_r$ and

$$\Gamma_\delta := \bigcup_{N \in \mathcal{N}_0} N^{-1}.$$

Recall the definition of N' for a nice rectangle N in the proof of Proposition 21. For any $N \in \mathcal{N}_0$, since $|\Gamma \cap N| \geq \delta |N|$ and $N^{-1} \subset N'$, by Proposition 21 we have

$$\min_{N^{-1}} u \geq \min_{N'} u \geq \gamma \min_{\Gamma} u.$$

Hence,

$$\min_{\Gamma_\delta} u \geq \gamma \min_{\Gamma} u.$$

Moreover, note that $\Gamma_\delta = Q_r$ when $\mathcal{N}_0 = \{Q_r\}$. Otherwise, if $\mathcal{N}_0 \neq \{Q_r\}$, we have

$$|\Gamma \cap N^{-1}| < \delta |N^{-1}| \quad \text{for all } N \in \mathcal{N}_0.$$

Therefore, if $\mathcal{N}_0 \neq \{Q_r\}$,

$$\begin{aligned} |\Gamma| = \Big| \bigcup_{N \in \mathcal{N}_0} (\Gamma \cap N) \Big| &\leq \Big| \bigcup_{N \in \mathcal{N}_0} (\Gamma \cap N^{-1}) \Big| \\ &< \sum_{N^{-1} : N \in \mathcal{N}_0} \delta |N^{-1}| = \delta |\Gamma_\delta|. \end{aligned}$$

Our proof is complete. □

Proof of Theorem 7:
We only consider the case when $\sigma < 1/7\sqrt{d}$.

For any $\Gamma \subset Q_{\sigma R}$, if $|Q_{\sigma R}| \leq \delta^{-s} |\Gamma|$ for some $s \in \mathbb{N}$, then we have

$$m := \min_{Q_{\sigma R}} u \geq \gamma^s \min_{\Gamma} u$$

by Lemma 22 and iteration. Hence for $t \geq 0$, putting $\Gamma^t := \{x \in Q_{\sigma R} : u(x) \geq t\}$, we get

$$m \geq \gamma^{\lceil \log_\delta (|\Gamma^t| / |Q_{\sigma R}|) \rceil} t \geq \left(\frac{|\Gamma^t|}{|Q_{\sigma R}|} \right)^{\log_\delta \gamma} \gamma t. \tag{51}$$

Note that $q := \log_\gamma \delta > 0$, since $\gamma, \delta \in (0,1)$. Therefore, for any $p \in (0,q)$,

$$\begin{aligned} \frac{1}{|Q_{\sigma R}|} \sum_{Q_{\sigma R}} u^p &= m^p + \frac{1}{|Q_{\sigma R}|} \sum_{Q_{\sigma R}} \int_m^\infty p t^{p-1} 1_{u \geq t} \, \mathrm{d}t \\ &= m^p + \int_m^\infty p t^{p-1} \frac{|\Gamma^t|}{|Q_{\sigma R}|} \, \mathrm{d}t \\ &\overset{(51)}{\leq} m^p + \int_m^\infty p t^{p-1} (\frac{m}{\gamma t})^q \, \mathrm{d}t \leq C m^p, \end{aligned}$$

where C depends on κ, d and $b_0 R$. Combining this and Theorem 19, the Harnack inequality for $\sigma < 1/7\sqrt{d}$ is proved.

Then case $\sigma \geq 1/7\sqrt{d}$ then follows by a chaining argument. □

References

[1] G. Ben Arous, Y. Hu, S. Olla, O. Zeitouni, *Einstein relation for biased random walk on Galton-Watson trees*, arXiv:1106.4387

[2] N. Berger, J.-D. Deuschel, *A quenched invariance principle for non-elliptic random walk in i.i.d. balanced random environment*, arXiv:1108.3995

[3] F. Comets, O. Zeitouni, *A law of large numbers for random walks in random mixing environments*, Ann. Probab. 32 (2004), no. 1B, 880-914.

[4] A. Einstein, *Investigations on the theory of the Brownian movement*, Edited with notes by R. Fürth. Translated by A. D. Cowper. Dover, New York, 1956.

[5] D. Gilbarg, N. Trudinger, *Elliptic partial differential equations of second order*, Second edition. Springer-Verlag, Berlin, 1983.

[6] N. Gantert, P. Mathieu, A. Piatnitski, *Einstein relation for reversible diffusions in random environment*, Comm. Pure Appl. Math. (2012) 65: 187-228.

[7] X. Guo, O. Zeitouni, *Quenched invariance principle for random walks in balanced random environment*, Probab. Theory Relat. Fields (2012) 152:207-230.

[8] T. Komorowsky, S. Olla, *Einstein relation for random walks in random environment*, Stochastic Process. Appl. 115 (2005), no. 8, 1279-1301.

[9] H. J. Kuo, N. S. Trudinger, *Linear elliptic difference inequalities with random coefficients*, Math. Comp. 55 (1990), 37–53.

[10] H. J. Kuo, N. S. Trudinger, *Positive difference operators on general meshes*, Duke Math. J. 83 (1996), 415–433.

[11] G. Lawler, *Weak convergence of a random walk in a random environment*, Comm. Math. Phys. 87 (1982/83), no. 1, 81-87.

[12] G. Lawler, *Estimates for differences and Harnack inequality for difference operators coming from random walks with symmetric, spatially inhomogeneous, increments*, Proc. London Math. Soc. (3) 63 (1991), no. 3, 552-568.

[13] J. Lebowitz, H. Rost, *The Einstein relation for the displacement of a test particle in a random environment*, Stochastic Process. Appl. 54 (1994), no. 2, 183-196.

[14] L. Shen, *On ballistic diffusions in random environment*, Ann. Inst. H. Poincaré Probab.Statist. 39(2003),no.5, 839-876.

[15] A.S. Sznitman, M. Zerner, *A law of large numbers for random walks in random environment*, Ann. Probab. 27 (1999), no. 4, 1851-1869.